

### Indexing Metadata/Description

- › **Title/condition:** Acoustic Neuroma: Physical Therapy
- › **Synonyms:** Neuroma, acoustic; vestibular schwannoma; acoustic schwannoma; acoustic neurinoma; cerebellopontine angle tumor; neurilemmoma; vestibular neuroma
- › **Anatomical location/body part affected:** Schwann cells surrounding the 8th cranial nerve, usually within the auditory canal
- › **Area(s) of specialty:** Neurological Rehabilitation
- › **Description:** <sup>(1,2,6)</sup>
  - An acoustic neuroma is a slow-growing, benign, intracranial, extra-axial tumor composed of Schwann cells that originates in the nerve sheath of the vestibulocochlear nerve, and usually on the superior vestibular portion. It usually arises in the internal auditory canal near the cerebellopontine angle
  - Accounts for 8 to 10% of all intracranial tumors and more than 90% of all cerebellopontine angle tumors
  - Patients typically present with hearing loss, tinnitus, and balance dysfunction, and might require physical therapy to address impairments in balance, gaze stability, functional mobility, and ability to ambulate, as well as facial weakness
- › **ICD-10 codes**
  - D33.3 Benign neoplasm of cranial nerves
  - R42 Dizziness and giddiness
  - R26.2 Difficulty in walking, not elsewhere classified
  - R26.81 Unsteadiness on feet
  - R26.89 Other abnormalities of gait and mobility

(ICD codes are provided for the reader's reference, not for billing purposes)

- › **Reimbursement:** No specific issues or information regarding reimbursement has been identified
- › **Presentation/signs and symptoms**
  - The majority of acoustic neuromas present unilaterally (95%)<sup>(2)</sup>
  - Bilateral presentation is much less common (5%) and is usually due to neurofibromatosis type 2 (NF2)<sup>(1)</sup>
  - Tumors associated with NF2 typically occur before the age of 30,<sup>(2)</sup> while those not associated with NF2 tend to present in the fifth and sixth decades<sup>(31)</sup>
  - Hearing loss is the most common symptom,<sup>(18,37)</sup> typically gradual onset of high-frequency sensorineural hearing loss,<sup>(4,6)</sup> with deterioration of speech discrimination exceeding that predicted by the degree of pure tone loss<sup>(1,4)</sup>
  - Vestibular dysfunction more commonly presents as continuous disequilibrium than episodic vertigo<sup>(1)</sup>
  - Tinnitus<sup>(4,6,37)</sup>
    - Occurs in approximately 63–75% of patients<sup>(12)</sup>
    - For detailed information, see Clinical Review... *Tinnitus*; CINAHL Topic ID Number: T709236

#### Author

Diane Matlick, PT  
Cinahl Information Systems, Glendale, CA

#### Reviewers

Amy Lombara, PT, DPT  
Cinahl Information Systems, Glendale, CA

Ellenore Palmer, BScPT, MSc  
Cinahl Information Systems, Glendale, CA

Rehabilitation Operations Council  
Glendale Adventist Medical Center,  
Glendale, CA

#### Editor

Sharon Richman, DHSc, MSPT, PT  
Cinahl Information Systems, Glendale, CA

January 1, 2021

- Dizziness<sup>(1,18,19)</sup>
  - Dizziness is the only symptom that seems to decrease as the size of the tumor increases<sup>(18)</sup>
- Unsteadiness/disequilibrium<sup>(1,5,37)</sup>
  - Researchers in Uruguay who conducted an RCT found that patients who were still unsteady one year after surgery to completely remove an acoustic neuroma had alterations in their response to gravitational vertical, as measured by the head-tilt response test (a test in which the patient wears goggles and has to tilt his or her head in order to keep a white bar aligned with his or her gravitational vertical)<sup>(5)</sup>
- Vertigo<sup>(1)</sup>
- Less common symptoms that might occur as tumor progresses
  - Headaches<sup>(2)</sup>
  - Diplopia<sup>(8)</sup>
  - Loss of coordination/ataxia<sup>(2)</sup>
  - Facial weakness<sup>(37)</sup>
  - Facial numbness<sup>(6)</sup>
  - Trigeminal nerve dysfunction<sup>(2)</sup>
  - Pyramidal weakness<sup>(37)</sup>
  - Nystagmus<sup>(37)</sup>

## Causes, Pathogenesis, & Risk Factors

### › Causes

- Atypical proliferation of Schwann cells
  - Idiopathic – unknown cause, usually unilateral<sup>(2)</sup>
  - Genetic mutation–manifestation of autosomal dominant disorder NF2. NF2 is caused by a mutation on the NF2 gene on the long arm of chromosome 22, and is often bilateral<sup>(2,15,16)</sup>
  - Recent studies have also identified increased risk of schwannomatosis, a member of the neurofibromatosis family of neurogenetic disorders, with mutation of the gene LZTR1<sup>(15,16)</sup>
    - Although schwannomas with this disease tend to be non-vestibular, cases of unilateral acoustic neuromas are observed<sup>(15)</sup>

### › Pathogenesis

- Benign, slowly growing encapsulated tumor composed of Schwann cells located around superior or inferior vestibular nerves that might lead to cochlear nerve dysfunction (due to compression or stretching), internal auditory artery occlusion, facial and trigeminal nerve dysfunction, increased intracranial pressure (ICP), or brainstem and/or cerebellar compression<sup>(37)</sup>
  - Compression of the facial and vestibulocochlear nerves when located in the internal acoustic canal<sup>(2)</sup>
  - Compression of the brain stem, 4<sup>th</sup> ventricle, and trigeminal nerve when located at the cerebellopontine angle<sup>(2)</sup>
  - The glossopharyngeal and vagus nerve are less commonly involved<sup>(31)</sup>
- The NF2 gene produces merlin, a tumor suppressor; therefore, deficiency in NF2 genes leads to the tumor development<sup>(2)</sup>

### › Risk factors

- Environmental exposure to high-dose ionizing radiation<sup>(11,37)</sup>
- NF2
  - Five percent of cases are due to NF2<sup>(1,11)</sup>
- Exposure to loud noise<sup>(11,35)</sup>
  - Results from studies are conflicting<sup>(11)</sup>
    - Researchers from Sweden who conducted a study in 2014 found no statistically significant association between acoustic neuroma and persistent occupational noise exposure, with or without hearing protection, but they did find a statistically

significant association between leisure-time exposure to loud noise, although they could not rule out a recall bias (i.e., the inability of the participants to correctly remember past events) as an alternative explanation<sup>(11)</sup>

- Researchers from Denmark who conducted a case-controlled study found no association between acoustic neuroma and residential traffic noise<sup>(35)</sup>
- Mobile phone use<sup>(7,10,33)</sup>
  - In studies in Denmark, the United Kingdom, Canada, and Sweden, researchers have found no increase in likelihood of acoustic neuroma with long-term mobile phone use. Most recently, researchers from Canada re-analyzed a prior study and did not find significant evidence of an increased risk for acoustic neuroma related to mobile phone use

## Overall Contraindications/Precautions

- › Any worsening of symptoms (e.g., impaired balance, gait, mental status) or appearance of new symptoms (e.g., headache, dizziness) should be immediately reported to the referring physician
- › Elevated ICP may present with vomiting, fever, and visual changes<sup>(31)</sup>
- › Examination and treatment should be performed by a trained vestibular therapist
- › See specific **Contraindications/precautions** under **Assessment/Plan of Care**

## Examination

### › History

#### • History of present illness/injury

##### –Mechanism of injury or etiology of illness

- When did symptoms develop? What were the initial symptoms? What are the primary complaints?
- When was the acoustic neuroma diagnosed?
- Does the patient have any of the risk factors?
- Does the patient have any hearing loss?<sup>(6)</sup>
  - Researchers who conducted a study in the United States found that workers at risk for hearing loss due to their jobs who undergo annual hearing exams as a part of a company-instituted hearing conservation program are more likely to have an acoustic neuroma diagnosed that otherwise would have gone undetected<sup>(6)</sup>
- What treatment did patient receive for the tumor?
- Were there any postoperative or posttreatment complications?
- How have patient's symptoms changed postoperatively or posttreatment?

##### –Course of treatment

- **Medical management:** Has the patient had surgery, radiosurgery, or radiotherapy, or is the acoustic neuroma under observation?
  - Observation
    - Patients with static acoustic neuromas should be advised that, although the tumor is not increasing in size, they still may be at increased risk for hearing loss in that ear<sup>(14)</sup>
    - Researchers from the Netherlands found increased quality of life in patients with a small acoustic neuroma who were managed with observation versus active treatment<sup>(34)</sup>
      - The study included 1,208 patients; the Penn Acoustic Neuroma Quality of Life questionnaire (PANQOL) was used for assessment
  - Microsurgery
    - Involves resection of the tumor; has the lowest rate of recurrence (up to 97.5% have complete tumor removal)<sup>(2)</sup>
    - Three standard approaches<sup>(2)</sup>
      - Retromastoid/retrosigmoid
      - Middle cranial fossa
      - Translabyrinthine
    - Staged resection<sup>(26)</sup>
      - Occasionally, during unfavorable cases involving large tumors, surgery has to be suspended to avoid risk to patient life. Additional surgeries are then necessary

- Stereotactic radiosurgery
  - Uses image-guided radiation directed right at the lesion; leads to disruption in cancer-cell DNA and, ultimately, cell death<sup>(21)</sup>
  - Delivery methods are gamma knife, linear accelerators, and proton beam and heavy-charged particles<sup>(21)</sup>
  - Often used in patients who are at high risk for complications associated with anesthesia<sup>(21)</sup>
  - Gamma knife single-dose stereotactic radiosurgery<sup>(2)</sup>
    - Performed on outpatient basis; indicated for smaller tumors (< 3 cm) or for patients with contraindications to microsurgery
    - Better tumor control rates than conservative management and fewer complications than high-dose radiation, but may not be as effective as high-dose
    - Complications include trigeminal or facial nerve damage and hydrocephalus
- Radiotherapy
  - Stereotactic radiation therapy
    - Typically used when the tumor is of substantial size and resection is not an option
    - Delivers higher dose of radiation to tumor and less to surrounding tissue
    - Requires multiple treatments
  - Hypofractionated stereotactic radiation treatment<sup>(9)</sup>
    - A more shortened course of therapy<sup>(9)</sup>
    - Preferable because it is more comfortable for the patient than surgery or conventional regimens but with comparable results, according to researchers in Austria<sup>(9)</sup>
- Chemotherapy has not been adequately investigated<sup>(2)</sup>
- Treatment for hearing loss<sup>(27,32)</sup>
  - Nonsurgical
    - Contralateral routing of signal hearing aid
    - Bone conduction devices
  - Surgical
    - Surgically implanted bone conduction devices or cochlear implants
      - Can be implanted at the time of tumor removal
      - Cochlear implants can only be used when the cochlear portion of the nerve is preserved<sup>(32)</sup>
- Treatment for facial nerve disruption<sup>(29)</sup>
  - Facial nerve reconstruction with graft
- **Medications for current illness/injury**
  - As the molecular mechanisms behind the genesis of these tumors are better understood, novel therapies are emerging that might reduce mortality and morbidity while preserving function<sup>(1,2,20)</sup>
  - Bevacizumab (vascular endothelial growth factor blocker) has shown promise for treatment of tumors in patients with NF2<sup>(1,31)</sup>
    - Has been shown to improve hearing and reduce volume of tumors<sup>(31)</sup>
  - Researchers in the United States studying salicylates report that they are a promising pharmacotherapy, as COX-2 is a key modulator in acoustic neuroma cell proliferation<sup>(20)</sup>
- **Diagnostic tests**
  - MRI
    - MRI should typically be the first diagnostic test performed when acoustic neuroma is suspected; using a noncontrast MRI first is more cost-effective and is generally as useful as an MRI with contrast<sup>(1)</sup>
    - MRI with gadolinium is the gold standard test and has 100% specificity<sup>(2)</sup>
    - Noncontrast T2-weighted fast-spin echo MRI is less expensive and has 98% specificity<sup>(2)</sup>
  - CT scan can have a high false-negative rate (as high as 37%)<sup>(2)</sup>
  - Audiometry<sup>(6)</sup>

- Stacked auditory brainstem response (ABR)<sup>(2)</sup>
  - ABR is highly sensitive and less expensive than MRI but its sensitivity is lower for smaller tumors<sup>(18)</sup>
- Electronystagmogram (ENG)
  - Caloric testing (part of the ENG) is the introduction of hot and cold air or water into the ear to test the vestibular ocular reflex (VOR). Typically it reveals a hypofunction on the affected side<sup>(30)</sup>
- Distortion-product otoacoustic emission tests can determine if the tumor is on the cochlear or vestibular portion of the nerve; retrocochlear hearing loss correlates with the cochlear portion<sup>(2)</sup>
- **Home remedies/alternative therapies:** Document any use of home remedies (e.g., ice or heating pack) or alternative therapies (e.g., acupuncture) and whether they help
- **Previous therapy:** Document whether patient has had occupational or physical therapy for this or other conditions and what specific treatments were helpful or not helpful
- **Aggravating/easing factors** (and length of time each item is performed before the symptoms come on or are eased): Are symptoms (e.g. tinnitus, dizziness, headache) associated with any specific activities or with head or body position changes? Does anything relieve or worsen symptoms?
- **Body chart:** Use body chart to document location and nature of symptoms
- **Nature of symptoms:** Document nature of symptoms (e.g., constant vs. intermittent, sharp, dull, aching, burning, numbness, tingling). Are symptoms such as dizziness, vertigo, imbalance, tinnitus, and/or headaches constant or intermittent? Ask patient to further describe (e.g. is tinnitus high pitch or low pitch? Is the patient dizzy or is the room spinning?)
- **Rating of symptoms:** Use a visual analog scale (VAS) or 0–10 scale to assess symptoms at their best, at their worst, and at the moment (specifically address if pain is present now and how much)
  - Activities-specific Balance Confidence (ABC) Scale: A 16-item scale in which patients rate confidence to perform specific activities
  - Visual Vertigo Analogue Scale (VVAS)
- **Pattern of symptoms:** Document changes in symptoms throughout the day and night, if any (a.m., mid-day, p.m., night); also document changes in symptoms due to weather or other external variables
- **Sleep disturbance:** Document number of wakings/night
- **Other symptoms:** Document other symptoms patient might be experiencing that could exacerbate the condition and/or symptoms that could be indicative of a need to refer to physician (e.g., symptoms of hydrocephalus such as worsening of balance, gait, or mental status)
- **Respiratory status:** Any respiratory complications during surgery? Does the patient require supplemental oxygen?
- **Barriers to learning**
  - Are there any barriers to learning? Yes \_\_ No \_\_
  - If Yes, describe \_\_\_\_\_
- **Medical history**
  - **Past medical history**
    - **Previous history of same/similar diagnosis**
      - Other vestibular disorders
      - CNS disorders
      - Hearing impairment
      - Any disease that affects the sensory systems (e.g., diabetes)
      - Vision problems (e.g., macular degeneration)
      - Migraine/other headache
    - **Comorbid diagnoses:** Ask patient about other problems, including diabetes, cancer, cardiovascular disease, psychiatric disorders, orthopedic disorders, and complications of pregnancy
      - Pregnancy may increase risk and accelerate growth of the tumor<sup>(2)</sup>
    - **Medications previously prescribed:** Obtain a comprehensive list of medications prescribed and/or being taken (including OTC drugs). Is the patient taking medications that affect the vestibular system, such as vestibular suppressants, sedatives, anticonvulsants, antihistamines, antipsychotics, or antidepressants?
    - **Other symptoms:** Ask patient about other symptoms he or she might be experiencing. Specifically ask about symptoms the patient might not realize are associated (e.g., fatigue, facial weakness, eye problems) or what he or she would like to speak about with a healthcare provider

- Researchers in the United Kingdom conducted an online survey (N = 480) to investigate the issues, needs, and concerns of patients undergoing treatment for acoustic neuroma<sup>(24)</sup>
  - The most frequently selected issues were tinnitus, fatigue, and dizziness
  - The most frequently selected fear was the tumor recurring
  - The most frequently selected concerns were facial appearance and head and neck pain

• **Social/occupational history**

- **Patient's goals:** Document what the patient hopes to accomplish with therapy and in general
- **Vocation/avocation and associated repetitive behaviors, if any:** Does the patient participate in recreational or competitive sports? Does the patient work? What does his or her job require? Does the patient participate in any other activities?
- **Functional limitations/assistance with ADLs/adaptive equipment**
  - Does the patient report any preexisting functional limitations?
  - Does patient have and/or use adaptive equipment?
- **Living environment:** Inquire about stairs, number of floors in home, and with whom patient lives. Does the patient have a caregiver? Identify if there are barriers to independence in the home. Are any modifications necessary? Does the home seem safe based on the information provided by the patient?

› **Relevant tests and measures: (While tests and measures are listed in alphabetical order, sequencing should be appropriate to patient medical condition, functional status, and setting)**

- **Arousal, attention, cognition (including memory, problem solving):** Assess in all patients and routinely monitor for changes. Consider using the Mini-Mental State Exam (MMSE), and obtain all formal testing results done by other disciplines
- **Assistive and adaptive devices:** Assess need for adaptive equipment for all patients with imbalance or dizziness. Assess proper fit and use for all equipment patient already has
- **Balance:** Assess all patients statically and dynamically. If possible, test with platform posturography (EquiTest) or the Clinical Test of Sensory Integration on Balance (CTSIB) (also known as the Foam and Dome test), as well as with Romberg test. Utilize standardized tests such as Berg Balance Scale, Tinetti Balance Assessment, Functional Reach Test, and standing on one leg
- **Cardiorespiratory function and endurance:** Assess blood pressure in lying, sitting, and standing. Assess vital signs throughout treatment as indicated and appropriate
- **Cranial/peripheral nerve integrity**
  - Assess for involvement of cranial nerves
    - I – Olfactory (evaluate ability to smell)
    - II – Optic (evaluate visual acuity, visual fields, and ocular fundi)
    - II, III – Optic and oculomotor (evaluate pupillary reactions)
    - III, IV, VI – Oculomotor, trochlear, abducens (evaluate extraocular movements)
    - V – Trigeminal (evaluate corneal reflexes, facial sensation, and jaw movements)
    - VII – Facial (evaluate facial movements, including lifting eyebrows, smiling/frowning, showing teeth, filling cheeks with air, and shutting eyes tight)
    - VIII – Vestibulocochlear (evaluate ability to hear)
    - IX, X – Glossopharyngeal, vagus (evaluate swallowing and rise of the palate, gag reflex)
    - V, VII, X, XII – Trigeminal, facial, vagus, hypoglossal (evaluate voice and speech)
    - XI – Spinal accessory (evaluate neck and shoulder movements)
    - XII – Hypoglossal (evaluate tongue symmetry and position)
  - Cranial nerve VIII is commonly affected by an acoustic neuroma<sup>(37)</sup>
  - Patients who have had surgery to remove an acoustic neuroma might have facial nerve (CN VII) involvement
  - Involvement of other cranial nerves (IV, VI, IX, and X, XI, XII) depends on the size of the tumor<sup>(1,37)</sup>
- **Functional mobility** (including transfers, etc.): Assess safety with mobility and note any symptoms with position changes. Use objective measurements such as the FIM and the Timed Up & Go (TUG) test
- **Gait/locomotion:** Assess for any imbalance, sway, and/or veering
  - Perform thorough gait analysis
  - Utilize standardized tests such as Dynamic Gait Index (DGI) to assess safety with ambulation
  - Vestibular specific gait tests/analysis
    - Ambulation with head turns side to side and up and down

- Researchers from the United States found that 4–8 weeks after resection for acoustic neuroma, patients still had deficits in head-trunk kinematics (i.e., turning head on body) during gait tasks<sup>(36)</sup>
- Walking with eyes closed
- **Joint integrity and mobility:** Assess cervical spine. If there is a history of pain or dysfunction, a thorough exam should be performed
- **Muscle strength:** Assessment of upper and lower extremities. If patient has facial weakness, assess strength of facial movements as well
  - For evaluating facial muscle weakness
    - Sunnybrook Facial Grading System
    - Facial Clinimetric Evaluation (FaCE) Scale<sup>(23)</sup>
- **Oral mechanism exam and related tests:** Perform oral mechanism exam if facial weakness is present
- **Range of motion:** Assess all motions in the cervical spine. Also assess hips and ankles, as normal ROM allows for optimal balance reactions
- **Self-care/activities of daily living** (objective testing): Assess safety in ADLs for patients with dizziness or imbalance; can use the Barthel Index
- **Special tests specific to diagnosis**
  - Gaze stability
    - Testing includes the following:
      - Spontaneous nystagmus
      - Gaze-evoked nystagmus
      - Rapid head thrust<sup>(30)</sup>
    - VOR (horizontal [HVOR], vertical [VVOR])
      - Head turns per minute while focused on object
      - 15° to each side
      - Normal is 120 head turns per minute
    - Static vs. dynamic visual acuity
    - Post head-shaking-nystagmus<sup>(30)</sup>
    - Vibration-induced nystagmus<sup>(30)</sup>
      - Hyperventilation test.<sup>(30)</sup> Researchers in Italy studied the hyperventilation test and found that hyperventilating for 60 seconds induced a nystagmus in 40/45 (89%) patients with unilateral acoustic neuroma, which is indicative of a unilateral vestibular deficit<sup>(30)</sup>
  - Use the Dix-Hallpike test to assess for benign paroxysmal positional vertigo (BPPV), which causes true vertigo (room spinning) for seconds to a minute when the patient lies down, rolls over, and looks up and/or down. For detailed information, see Clinical Review... *Benign Paroxysmal Positional Vertigo*; CINAHL Topic ID Number: T708435
    - In order to perform the test, begin with patient in long sitting position, rotate head 45° to one side and have patient lie back with head hanging off edge of bed. Wait 45 seconds and sit back up. Repeat on other side. Positive test indicated by patient's subjective report of the room spinning and therapist's objective report of rotary nystagmus. Horizontal nystagmus is possible but less likely to result in a positive response, and indicates horizontal canal BPPV
  - Motion sensitivity: Assess for 16 specific positions. Patient reports and therapist notes severity and duration of symptoms
  - Dizziness Handicap Inventory (DHI): 25-item questionnaire that evaluates how dizziness affects the patient's quality of life, including emotional, functional, and physical domains
  - Tinnitus Functional Index (TFI)<sup>(23)</sup>
  - Quality of life<sup>(25)</sup>
    - PANQOL
      - Minimally Clinically Important Difference (MCID) is 11
    - SF-36
      - MCID for mental health component is 7 and for physical health component is 8

## Assessment/Plan of Care

### › **Contraindications/precautions**

- When possible, treatment should be provided by a trained vestibular therapist

- **Patients with this diagnosis might be at risk for falls; follow facility protocols for fall prevention and post fall-prevention instructions at bedside, if inpatient. Inform patient and family/caregivers of the potential for falls and educate about fall-prevention strategies. Discharge criteria should include independence with fall-prevention strategies**

› **Diagnosis/need for treatment**

- Patients with acoustic neuroma benefit from physical therapy to treat dizziness, imbalance, gait abnormality, and facial weakness

› **Rule out**

- Labyrinthitis
- Vestibular neuritis
- BPPV<sup>(31)</sup>
- Migraine-associated dizziness
- Ménière's disease<sup>(31)</sup>
- Perilymphatic fistula
- Microvascular compression
- Chiari malformation
- Gentamicin toxicity<sup>(31)</sup>
- Multiple sclerosis
- Hearing loss of different etiology<sup>(2)</sup>
  - Presbycusis (age-related hearing loss)
  - Middle ear effusion
  - Infection
  - Wax
  - Cholesteatoma
  - Tympanic membrane rupture
- Stroke<sup>(31)</sup>
- Brain injury
- Other cerebellopontine angle tumors (e.g. meningiomas, epidermoid tumors)<sup>(19)</sup>

› **Prognosis**

- Without treatment, approximately half of acoustic neuromas shrink or do not grow further<sup>(2)</sup>
- Conservative therapy is more likely to preserve hearing than radiotherapy or surgery<sup>(2)</sup>
- Researchers in the Czech Republic who analyzed the effect of microsurgery on hearing and tinnitus found that the main prognostic indicators of hearing preservation were grade/size of tumor, preoperative hearing level, intraoperative neuromonitoring, tumor consistency, and adhesion to neurovascular structures. The main prognostic indicators of tinnitus elimination were preservation of useful hearing and neurectomy of the eighth cranial nerve. The main prognostic indicators for persistence and new onset of tinnitus were the preservation of cochlear nerve but loss of preoperative hearing<sup>(13)</sup>
- Researchers in the United States found that 50–70% of patients maintained serviceable hearing after surgery using middle cranial fossa approach and, of those, 75% maintained serviceable hearing after 5 years<sup>(28)</sup>
- Researchers who conducted a study in the United States found similar quality of life outcomes for patients regardless of management (conservative, gamma knife, or surgery)<sup>(3)</sup>
  - Main outcome used was PANQOL scores

› **Referral to other disciplines**

- Social work for needs at home
- Psychology for anxiety, depression
- Speech therapy for patients with impaired speech or dysphagia
- Occupational therapy for ADL training
- Physician for red flags, suspicion of hydrocephalus
- Audiologist for hearing testing, vestibular testing, hearing aids

› **Other considerations**

- Patients might have the following problems in association with acoustic neuroma



–Hydrocephalus<sup>(2)</sup>

- Symptoms include headache, ataxia, worsening of balance or gait, blurred or double vision, urinary incontinence, decline of mental status

–Brainstem compression<sup>(2)</sup>

–Cerebellar tonsil herniation<sup>(2)</sup>

• Surgical complications

–Possible complications include the following:

- Headaches

- Hearing loss

- Although rare, can affect contralateral ear as well<sup>(12)</sup>

- Facial paresis<sup>(37)</sup>

- Cerebellar injury

- Damage to the anterior or posterior intracerebral arteries<sup>(37)</sup>

- CSF leakage can lead to meningitis and abscess formation in the brain<sup>(17)</sup>

- Air embolism

- Intracranial hemorrhage

- Impaired vestibular compensation<sup>(21)</sup>

- Meningitis<sup>(21)</sup>

- Mortality<sup>(21)</sup>

› **Treatment summary**

• There is limited research regarding physical therapy for patients with acoustic neuroma

–Limitations of studies include small sample sizes, lack of randomization, and lack of controls or confounding variables

<b>Problem</b>	<b>Goal</b>	<b>Intervention</b>	<b>Expected Progression</b>	<b>Home Program</b>
Impaired functional mobility due to balance, flexibility, and strength impairments	Improved balance, flexibility, and strength resulting in improved functional mobility	<b><u>Therapeutic exercise</u></b>  General strengthening and/or stretching exercises if indicated to maximize strength and flexibility to allow for optimal balance reactions  Romberg exercises, static and dynamic balance exercises that vary somatosensory, visual, and vestibular conditions	Strengthening exercises can be progressed with resistive equipment such as weights or from supine to sitting to standing  Balance activities can be progressed by narrowing base of support and/or having patient stand on one limb or unstable surfaces. Also can add head movements and cognitive activities	Provide patient/ family with written instructions and diagrams to safely perform exercises at home
Decreased ability to ambulate safely	Improved safety with ambulation	<b><u>Gait training</u></b>  Gait training, with assistive device if necessary	Wean off assistive device. Add head turns, obstacles, and cognitive tasks, vary surface, narrow base of support, and have patient close eyes	Provide patient/ family with written instructions and diagrams to perform safely at home

<p>Decreased safety with increased risk for falls while performing ADLs and functional mobility</p>	<p>Improve safety with ADLs performance and functional mobility</p>	<p><b><u>Functional and safety training</u></b></p> <p>Educate patient/ family regarding safe techniques for transfers, gait, and ADLs. Check that patient is using assistive devices and adaptive equipment correctly. Home modifications (e.g., remove throw rugs)</p>	<p>Provide less assistance and feedback as patient improves</p>	<p>Provide patient/ family with written instructions and diagrams regarding fall prevention techniques and home modifications</p>
<p>Decreased gaze stability, dizziness</p>	<p>Increased gaze stability, decreased dizziness</p>	<p><b><u>Vestibular exercises</u></b></p> <p>VOR X1 viewing: instruct patient to view a stationary object while turning his or her head side to side (HVOR) and up and down (VVOR) for 1–2 minutes while the object stays in perfect focus</p> <p>VOR X2 viewing: the patient views a target moving in the opposite direction of head rotation</p>	<p>As patient improves, he or she should be able to turn head faster with the object still staying perfectly clear. Adaptations to the vestibular exercises should be made as appropriate for each patient; take into consideration the environment in which patient performs work/ leisure or ADL activity</p> <p>Can progress to include different backgrounds (e.g., checkerboard) or perform exercises outdoors</p> <p>Incorporate VOR exercises with balance activities</p> <p>Normal rate is 120 head turns/minute</p>	<p>Perform exercises for 1–2 minutes, 2–3 times per day as tolerated</p>

Weakness of facial muscles	Increase strength of facial muscles	<p><b><u>Therapeutic exercise</u></b></p> <p>Facial muscle strengthening exercises might include exercises based on proprioceptive neuromuscular facilitation (PNF). PNF utilizes traction, contralateral contraction, manual contact, maximal resistance, verbal input, and stretching to increase motor control and strength</p>	Facial muscle strengthening can be progressed from passive to active assisted to active to resisted exercise	Provide patient/family with written instructions and diagrams
----------------------------	-------------------------------------	--	--	---

## Desired Outcomes/Outcome Measures

› Desired outcomes and associated measures

- Improved balance
  - Platform posturography (EquiTest)
  - Berg Balance Scale
  - Tinetti Balance Assessment
  - Functional Reach Test
  - Clinical Test of Sensory Integration on Balance
  - Standing on one leg
  - Romberg test
- Improved dizziness and vertigo
  - VVAS
  - UCLA Dizziness Questionnaire
  - Motion sensitivity test
- Improved safety and independence with ADLs and gait
  - DGI
  - TUG test
  - FIM
  - ABC Scale
  - DHI
  - Functional Checklist
- Increased gaze stability
  - VOR speed (head turns per minute)
- Increased strength of facial muscles
  - Sunnybrook Facial Grading System
  - FaCE Scale<sup>(22)</sup>
- Improved tinnitus
  - TFI
- Patient satisfaction/quality of life
  - PANQOL
  - SF-36

## Maintenance or Prevention

- › Patient is advised to follow home program and follow up with physician as indicated
- › Implementation of all prevention strategies and home modifications

## Patient Education

- › Mayo Clinic website: <https://www.mayoclinic.org/diseases-conditions/acoustic-neuroma/symptoms-causes/syc-20356127>
- › National Institute on Deafness and Other Communication Disorders website: <https://www.nidcd.nih.gov/health/vestibular-schwannoma-acoustic-neuroma-and-neurofibromatosis>
- › American Hearing Research Foundation website: <https://www.american-hearing.org/understanding-hearing-balance/>

## Note

- › Recent review of the literature has found no updated research evidence on this topic since previous publication on January 4, 2019

## Coding Matrix

References are rated using the following codes, listed in order of strength:

<b>M</b> Published meta-analysis	<b>RV</b> Published review of the literature	<b>PP</b> Policies, procedures, protocols
<b>SR</b> Published systematic or integrative literature review	<b>RU</b> Published research utilization report	<b>X</b> Practice exemplars, stories, opinions
<b>RCT</b> Published research (randomized controlled trial)	<b>QI</b> Published quality improvement report	<b>GI</b> General or background information/texts/reports
<b>R</b> Published research (not randomized controlled trial)	<b>L</b> Legislation	<b>U</b> Unpublished research, reviews, poster presentations or other such materials
<b>C</b> Case histories, case studies	<b>PGR</b> Published government report	<b>CP</b> Conference proceedings, abstracts, presentation
<b>G</b> Published guidelines	<b>PFR</b> Published funded report	

## References

- Lustig LR, Schindler JS. Ear, nose, & throat disorders. In: Papadakis MA, McPhee SJ, eds. *Current Medical Diagnosis & Treatment 2018*. 57th ed. New York, NY: McGraw-Hill Medical; 2018:218. **(GI)**
- Cohn J, Quercetti K, Ghaderi M. Acoustic neuroma. Domino FJ, ed. *The 5-Minute Clinical Consult Standard 2016*. 24th ed. Philadelphia, PA: Wolters Kluwer Health; 2015. **(GI)**
- McLaughlin EJ, Bigelow DC, Lee JY, Ruckenstein MJ. Quality of life in acoustic neuroma patients. *Otol Neurotol*. 2015;36(4):653-656. doi:10.1097/MAO.0000000000000674. **(R)**
- Lee SH, Choi SK, Lim YJ, et al. Otologic manifestations of acoustic neuroma. *Acta Otolaryngol*. 2015;135(2):140-146. doi:10.3109/00016489.2014.952334. **(R)**
- Suarez H, Ferreira E, Arocena S, et al. Chronic balance disorders after acoustic neuroma surgery: Assessment of gravitational vertical perception. *Acta Otolaryngol*. 2015;135(4):348-353. doi:10.3109/00016489.2014.974287. **(R)**
- Taiwo O, Galusha D, Tessier-Sherman B, et al. Acoustic neuroma: Potential risk factors and audiometric surveillance in the aluminum industry. *Occup Environ Med*. 2014;71(9):624-628. doi:10.1136/oemed-2014-102094. **(C)**
- Bensen VS, Pirie K, Schuz J, Reeves GK, Beral V, Green J. Authors' response to: The case of acoustic neuroma: Comment on mobile phone use and risk of brain neoplasms and other cancers. *Int J Epidemiol*. 2014;43(1):275. doi:10.1093/ije/dyt186. **(X)**
- Zhidan L, Jun H. Electroacupuncture therapy for abducent palsy after acoustic neuroma surgery. *Acupunct Med*. 2015;33(2):168-169. doi:10.1136/acupmed-2014-010699. **(C)**
- Kranzinger M, Zehentmayr F, Fastner G, et al. Hypofractionated stereotactic radiotherapy of acoustic neuroma: Volume changes and hearing results after 89-month median follow-up. *Strahlenther Onkol*. 2014;190(9):798-805. doi:10.1007/s00066-014-0630-4. **(R)**
- Pettersson D, Mathiesen T, Prochazka M, et al. Long-term mobile phone use and acoustic neuroma risk. *Epidemiology*. 2014;25(2):233-241. doi:10.1097/EDE.0000000000000058. **(R)**
- Fisher JL, Pettersson D, Palmisano S, et al. Loud noise exposure and acoustic neuroma. *Am J Epidemiol*. 2014;180(1):58-67. doi:10.1093/aje/kwu081. **(R)**
- Deeb RH, Rock JP, Seidman MD. Contralateral hearing loss after vestibular schwannoma excision: A rare complication of neurotologic surgery. *Ear Nose Throat J*. 2015;94(1):28-31. doi:N/A. **(C)**
- Chovanec M, Zverina E, Profant O, et al. Does attempt at hearing preservation microsurgery of vestibular schwannoma affect postoperative tinnitus? *BioMed Res Int*. 2015;1-9. doi:10.1155/2015/783169. **(R)**
- Patel NB, Nieman CL, Redleaf M. Hearing in static unilateral vestibular schwannoma declines more than in the contralateral ear. *Ann Otol Rhinol Laryngol*. 2015;124(6):490-4. doi:10.1177/0003489414566181. **(R)**
- Smith MJ, Isidor B, Beetz C, et al. Mutations in LZTR1 add to the complex heterogeneity of schwannomatosis. *Neurology*. 2015;84(2):141-7. doi:10.1212/WNL.0000000000001129. **(R)**
- Piotrowski A, Xie J, Liu YF, et al. Germline loss-of-function mutations in LZTR1 predispose to an inherited disorder of multiple schwannomas. *Nat Genet*. 2014;46(2):182-7. doi:10.1038/ng.2855. **(R)**
- Crowson MG, Cunningham CD, Moses H, Zomorodi AR, Kaylie DM. Preoperative lumbar drain use during acoustic neuroma surgery and effect on CSF leak incidence. *Ann Otol Rhinol Laryngol*. 2016;125(1):63-8. doi:10.1177/0003489415597917. **(R)**
- Lee SH, Choi SK, Lim YK, et al. Otologic manifestations of acoustic neuroma. *Acta Otolaryngol*. 2015;135(2):140-6. doi:10.3109/00016489.2014.952334. **(R)**
- Kim SH, Lee SH, Choi SK, Lim YJ, Na SY, Yeo SG. Audiologic evaluation of vestibular schwannoma and other cerebellopontine angle tumors. *Acta Otolaryngol*. 2016;136(2):149-53. doi:10.3109/00016489.2015.1100326. **(R)**
- Dilwali S, Kao SY, Fujita T, Landegger LD, Stankovic KM. Nonsteroidal anti-inflammatory medications are cytostatic against human vestibular schwannomas. *J Lab Clin Med*. 2015;166(1):1-11. doi:10.1016/j.trsl.2014.12.007. **(R)**
- Brathwaite J, Mendiratta P. Radio surgical ablation of acoustic neuromas: a unique treatment modality for an elderly woman. *J Am Geriatr Soc*. 2016;64(2):446-7. doi:10.1111/jgs.13979. **(C)**
- Leong SC, Lesser TH. A national survey of facial paralysis on the quality of life of patients with acoustic neuroma. *Otol Neurotol*. 2015;36(3):503-9. doi:10.1097/MAO.0000000000000428. **(R)**

23. Overdeest JB, Pross SE, Cheung SW. Tinnitus following treatment for sporadic acoustic neuroma. *Laryngoscope*. 2015. doi:10.1002/lary.25672. **(R)**
24. Leong SC, Lesser TH. A United Kingdom survey of concerns, needs, and priorities reported by patients diagnosed with acoustic neuroma. *Otol Neurotol*. 2015;36(3):486-90. doi:10.1097/MAO.0000000000000556. **(R)**
25. Carlson ML, Tveiten OV, Yost KJ, Lohse CM, Lund-Johnson M, Link ML. The minimally clinically important difference in vestibular schwannoma quality-of-life assessment: An important step beyond P <.05. *Otolaryngol Head Neck Surg*. 2015;153(2):202-8. doi:10.1177/0194599815585508. **(R)**
26. Kunimoto Y, Lauda L, Falcioni M, Taibah A, Hasegawa K, Sanna M. Staged resection for vestibular schwannoma. *Acta Otolaryngol*. 2015;135(9):895-900. doi:10.3109/00016489.2015.1040170. **(C)**
27. McRackan TR, Goddard JC, Wilkinson EP, Slattery WH, Brackman DE. Bone-anchored hearing device placement with translabyrinthine tumor removal. *Otolaryngol Head Neck Surg*. 2015;152(2):314-18. doi:10.1177/0194599814558038. **(R)**
28. Quist TS, Givens DJ, Gurgel RK, Chamoun R, Shelton C. Hearing preservation after middle fossa vestibular schwannoma removal: Are the results durable? *Otolaryngol Head Neck Surg*. 2015;153(4):706-11. doi:10.1177/0194599814567874. **(R)**
29. Ramos DS, Bonnard D, Franco-Vidal V, Liguoro D, Darrouzet V. Stitchless fibrin glue-aided facial nerve grafting after cerebellopontine angle schwannoma removal: technique and results in 15 cases. *Otol Neurotol*. 2015;36(3):498-502. doi:10.1097/MAO.0000000000000408. **(R)**
30. Califano L, Iorio G, Salafia F, Mazzone S, Califano M. Hyperventilation-induced nystagmus in patients with vestibular schwannoma. *Otol Neurotol*. 2015;36(2):303-306. doi:10.1097/MAO.0000000000000699. **(R)**
31. Bilodeau CC. Acoustic Neuroma. In: Ferri FF, ed. *2018 Ferri's Clinical Advisor*. Philadelphia, PA: Elsevier Inc; 2018:16. **(GI)**
32. DeHart AN, Broaddus WC, Coelho DH. Translabyrinthine vestibular schwannoma resection with simultaneous cochlear implantation. *Cochlear Implants Int*. 2017;18(5):278-284. doi:10.1080/14670100.2017.1337665. **(C)**
33. Momoli F, Siemiatycki J, McBride ML, et al. Probabilistic multiple-bias modeling applied to the canadian data from the interphone study of mobile phone use and risk of glioma, meningioma, acoustic neuroma, and parotid gland tumors. *Am J Epidemiol*. 2017;186(7):885-893. doi:10.1093/aje/kwx157. **(R)**
34. Soulier G, van Leeuwen BM, Putter H, et al. Quality of life in 807 patients with vestibular schwannoma: comparing treatment modalities. *Otolaryngol Head Neck Surg*. 2017;157(1):92-98. doi:10.1177/0194599817695800. **(R)**
35. Roswall N, Stangerup SE, Caye-Thomasen P, et al. Residential traffic noise exposure and vestibular schwannoma – a Danish case–control study. *Acta Oncol*. 2017;56(10):1310-1316. doi:10.1080/0284186X.2017.1337925. **(R)**
36. Paul SS, Dibble LE, Walther RG, Shelton C, Gurgel RIK, Lester ME. Characterization of head-trunk coordination deficits after unilateral vestibular hypofunction using wearable sensors. *JAMA Otolaryngol Head Neck Surg*. 2017;143(10):1008-1014. doi:10.1001/jamaoto.2017.1443. **(R)**
37. Greene J, Al-Dhahir MA. Acoustic neuroma (vestibular schwannoma) [Updated 2018 Nov 25]. StatPearls [Internet] Web site. <https://www.ncbi.nlm.nih.gov/books/NBK470177/>. Published 2019. **(GI)**